

# Treat More with the Only 300 mm DCB\*

Lutonix® Drug Coated Balloon family of PTA Catheters has the broadest size offering of all U.S. drug coated balloons (DCBs)\*, with sizes on both .018" and .035", indications in AV and femoropopliteal lesions, and in lengths up to 300 mm on the .018" platform. Now with the longest DCB length available, Lutonix® 018 DCB offers a comprehensive size matrix to treat lesions without the need to exchange to an .035" guidewire.

Balloon Diameter (mm)	Balloon Length (mm)						
	40	60	80	100	150	220	<b>300</b>
4	4F	4F	4F	4F	4F	4F	<b>5F</b>
5	5F	5F	5F	5F	5F	5F	<b>5F</b>
6	5F	5F	5F	5F	5F	5F	<b>5F</b>
7	5F	5F	5F	5F	5F	5F	5F

40 60 80 100 150 220 300

## Lutonix® 018 Drug Coated Balloon PTA Catheter

\* As of January 2020; on the US market



Diameter (mm)	Length (mm)	Sheath Profile	100 cm Catheter Length
4	40	4F	<input type="checkbox"/> LX181004404F
	150	4F	<input type="checkbox"/> LX1810041504F
	220	4F	<input type="checkbox"/> LX1810042204F
	300	5F	<input type="checkbox"/> LX1810043005F
5	40	5F	<input type="checkbox"/> LX181005405F
	150	5F	<input type="checkbox"/> LX1810051505F
	220	5F	<input type="checkbox"/> LX1810052205F
	300	5F	<input type="checkbox"/> LX1810053005F
6	40	5F	<input type="checkbox"/> LX181006405F
	150	5F	<input type="checkbox"/> LX1810061505F
	220	5F	<input type="checkbox"/> LX1810062205F
	300	5F	<input type="checkbox"/> LX1810063005F
7	40	5F	<input type="checkbox"/> LX181007405F
	150	5F	<input type="checkbox"/> LX1810071505F

Diameter (mm)	Length (mm)	Sheath Profile	130 cm Catheter Length
4	40	4F	<input type="checkbox"/> LX181304404F
	60	4F	<input type="checkbox"/> LX181304604F
	80	4F	<input type="checkbox"/> LX181304804F
	100	4F	<input type="checkbox"/> LX1813041004F
	150	4F	<input type="checkbox"/> LX1813041504F
	220	4F	<input type="checkbox"/> LX1813042204F
	300	5F	<input type="checkbox"/> LX1813043005F
5	40	5F	<input type="checkbox"/> LX181305405F
	60	5F	<input type="checkbox"/> LX181305605F
	80	5F	<input type="checkbox"/> LX181305805F
	100	5F	<input type="checkbox"/> LX1813051005F
	150	5F	<input type="checkbox"/> LX1813051505F
	220	5F	<input type="checkbox"/> LX1813052205F
	300	5F	<input type="checkbox"/> LX1813053005F
6	40	5F	<input type="checkbox"/> LX181306405F
	60	5F	<input type="checkbox"/> LX181306605F
	80	5F	<input type="checkbox"/> LX181306805F
	100	5F	<input type="checkbox"/> LX1813061005F
	150	5F	<input type="checkbox"/> LX1813061505F
	220	5F	<input type="checkbox"/> LX1813062205F
	300	5F	<input type="checkbox"/> LX1813063005F
7	40	5F	<input type="checkbox"/> LX181307405F
	60	5F	<input type="checkbox"/> LX181307605F
	80	5F	<input type="checkbox"/> LX181307805F
	100	5F	<input type="checkbox"/> LX1813071005F
	150	5F	<input type="checkbox"/> LX1813071505F
	220	5F	<input type="checkbox"/> LX1813072205F

I authorize the purchase of these products.

\_\_\_\_\_

PHYSICIAN NAME

\_\_\_\_\_

PHYSICIAN SIGNATURE

\_\_\_\_\_

REPRESENTATIVE'S NAME

\_\_\_\_\_

CONTACT PHONE NO.

**Lutonix® 018 Drug Coated Balloon PTA Catheter**

**Indications for Use:** The Lutonix® 018 Drug Coated Balloon PTA catheter is indicated for percutaneous transluminal angioplasty, after appropriate vessel preparation, of de novo, restenotic, or in-stent restenotic lesions up to 300 mm in length in native superficial femoral or popliteal arteries with reference vessel diameters of 4-7 mm. The Lutonix® 018 Drug Coated Balloon PTA catheter is indicated for percutaneous transluminal angioplasty, after pre-dilatation, for treatment of stenotic lesions of dysfunctional native arteriovenous dialysis fistulae that are 4 mm to 7 mm in diameter and up to 80 mm in length.

**Contraindications:** The Lutonix® Catheter is contraindicated for use in: - Patients who cannot receive recommended anti-platelet and/or anticoagulant therapy (SFA). - Women who are breastfeeding, pregnant or are intending to become pregnant or men intending to father children over the next two years. It is unknown whether paclitaxel will be excreted in human milk and there is a potential for adverse reaction in nursing infants from paclitaxel exposure. - Patients judged to have a lesion that prevents complete inflation of an angioplasty balloon or proper placement of the delivery system.

**Warnings:** - A signal for increased risk of late mortality has been identified following the use of paclitaxel-coated balloons and paclitaxel-eluting stents for femoropopliteal arterial disease beginning approximately 2-3 years post-treatment compared with the use of non-drug coated devices. There is uncertainty regarding the magnitude and mechanism for the increased late mortality risk, including the impact of repeat paclitaxel device exposure. Inadequate information is available to evaluate the potential mortality risk associated with the use of paclitaxel-coated devices for the treatment of other diseases/conditions, including this device indicated for use in arteriovenous dialysis fistulae. Physicians should discuss this late mortality signal and the benefits and risks of available treatment options with their patients. - Contents supplied STERILE using ethylene oxide (EO) process. Do not use if sterile barrier is damaged or opened prior to intended use. - Do not use after "Use by" date. - Do not use if product damage is evident. - The Lutonix® Catheter is for use in one patient only; do not reuse in another patient, reprocess or resterilize. Risks of reuse in another patient, reprocessing, or resterilization include: - Compromising the structural integrity of the device and/or device failure which, in turn, may result in patient injury, illness or death. - Creating a risk of device contamination and/or patient infection or cross-infection, including, but not limited to, the transmission of infectious disease(s) from one patient to another. Contamination of the device may lead to patient injury, illness or death. - Do not exceed the Rated Burst Pressure (RBP) recommended for this device. Balloon rupture may occur if the RBP rating is exceeded. To prevent over-pressurization, use of a pressure monitoring device is recommended. - Use the recommended balloon inflation medium of contrast and sterile saline (≤50% contrast). Never use air or any gaseous medium to inflate the balloon as this may cause air emboli in case of balloon burst. - This product should not be used in patients with known hypersensitivity to paclitaxel or structurally related compounds, as this may cause allergic reaction difficulty in breathing, skin rash, muscle pain.

**Precautions:** **General precautions:** - The Lutonix® Catheter should only be used by physicians trained in percutaneous interventional procedures. - Consideration should be given to the risks and benefits of use in patients with a history of non-controllable allergies to contrast agents. - The safety and effectiveness of the Lutonix® Catheter have not been established for treatment in cerebral, carotid, coronary, or renal vasculature. - For SFA application, the safety and effectiveness of using more than four Lutonix drug coated balloons or a maximum drug coating quantity of approximately 15.1 mg paclitaxel in a patient has not been clinically evaluated. - For AV Fistula application, the safety and effectiveness of using multiple Lutonix drug coated balloons that deliver greater than 7.6 mg paclitaxel in a patient has not been clinically evaluated.

Potential adverse events which may be associated with a peripheral balloon dilatation procedure include, but are not limited to the following: - Additional intervention - Allergic reaction to drugs, excipients or contrast medium - Amputation/loss of limb (SFA) - Aneurysm or pseudoaneurysm - Arrhythmias - Embolization - Hematoma - Hemorrhage, including bleeding at the puncture site - Hypotension/hypertension - Inflammation - Loss of permanent access (AVF) - Occlusion - Pain or tenderness - Pneumothorax or hemothorax (SFA) - Sepsis/infection - Shock - Steal Syndrome (AVF) - Stroke - Thrombosis - Vessel dissection, perforation, rupture, or spasm

Although systemic effects are not anticipated, refer to the Physicians' Desk Reference for more information on the potential adverse events observed with paclitaxel.

Potential adverse events, not described in the above source, which may be unique to the paclitaxel drug coating include, but are not limited to the following: - Allergic/immunologic reaction to the drug coating (paclitaxel) - Alopecia - Anemia - Blood product - transfusion - Gastrointestinal symptoms - Hematologic dyscrasia (including leukopenia, neutropenia, thrombocytopenia) - Hepatic enzyme changes - Histologic changes in vessel wall, including inflammation, cellular damage, or necrosis - Myalgia/Arthralgia - Myelosuppression - Peripheral neuropathy

Please consult product labels and instructions for use for indications, contraindications, hazards, warnings and precautions. **Px only**